

# Relativistic heavy quarks on the lattice

Sinya Aoki<sup>a</sup>, Yoshinobu Kuramashi<sup>b</sup> and Shin-ichi Tominaga<sup>c</sup>

<sup>a</sup>Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan

<sup>b</sup> High Energy Accelerator Research Organization(KEK), Tsukuba, Ibaraki 305-0801, Japan

<sup>c</sup> Center for Computational Physics, University of Tsukuba, Tsukuba, Ibaraki 305-8577, Japan

We investigate feasibility of relativistic approaches to the heavy quark physics on both anisotropic and isotropic lattices. Our perturbative calculation reveals that the anisotropic lattice is not theoretically advantageous over the isotropic one to control  $m_Q a$  errors. We instead propose a new relativistic approach to handle heavy quarks on the isotropic lattice.

## 1. INTRODUCTION

Lattice QCD is expected to provide the opportunity of precise evaluation for weak matrix elements associated with  $B$  mesons from first principles. The main obstacle on the way is the systematic errors originating from large  $m_b a$  errors. At present this difficulty is avoided in most cases by employing the nonrelativistic approaches[1,2]. On the other hand, the use of anisotropic lattice[3] has a fascinating feature to allow us to take the continuum limit. However, there exists a theoretical concern whether or not  $m_Q a_s$  errors could revive perturbatively or nonperturbatively even after they are removed classically. Unfortunately, our one-loop calculation of the quark self energy strongly suggests that radiative corrections allow the revival of  $m_Q a_s$  errors.

Having found that the anisotropic lattice is not theoretically advantageous over the isotropic one, we propose a new relativistic way to deal with the heavy quarks on the isotropic lattice. Cutoff effects in the heavy quark system are discussed by applying the on-shell improvement program[4,5] to the finite  $m_Q a$  case. We show that a proper adjustment of four parameters in the quark action reduces the remaining cutoff effects to be  $O((a\Lambda_{\text{QCD}})^2)$ . We also demonstrate a determination of the four parameters at the tree-level from the on-shell quark-quark scattering amplitude.

In this report we present the salient points in our work[6].

## 2. ANISOTROPIC LATTICE

In order to obtain a generic form of the quark action on the anisotropic lattice, let us make the operator analysis according to the on-shell improvement program[4,5]. Under the requirement of various symmetries on the lattice, we find the following set of operators are allowed up to dimension five:

$$\text{dim.3 : } \quad \mathcal{O}_3(x) = \bar{q}(x)q(x), \quad (1)$$

$$\text{dim.4 : } \quad \mathcal{O}_{4a}(x) = \bar{q}(x)\gamma_0 D_0 q(x), \quad (2)$$

$$\mathcal{O}_{4b}(x) = \sum_i \bar{q}(x)\gamma_i D_i q(x), \quad (3)$$

$$\text{dim.5 : } \quad \mathcal{O}_{5a}(x) = \bar{q}(x)D_0^2 q(x), \quad (4)$$

$$\mathcal{O}_{5b}(x) = \sum_i \bar{q}(x)D_i^2 q(x), \quad (5)$$

$$\mathcal{O}_{5c}(x) = i\sum_i \bar{q}(x)\sigma_{0i}F_{0i}q(x), \quad (6)$$

$$\mathcal{O}_{5d}(x) = i\sum_{i,j} \bar{q}(x)\sigma_{ij}F_{ij}q(x), \quad (7)$$

$$\mathcal{O}_{5e}(x) = \sum_i \bar{q}(x)[\gamma_0 D_0, \gamma_i D_i]q(x), \quad (8)$$

where the subscript 0 denotes the time component, while  $i, j = 1, 2, 3$  space components. Two degree of freedom in the eight coefficients are absorbed in the renormalization of the quark mass  $Z_m$  and the wave function  $Z_q$ . We also find  $\mathcal{O}_{5a}$  and  $\mathcal{O}_{5e}$  are related to other operators by using the eq. of motion and hence they are redundant. The remaining four parameters have to be tuned to remove  $O(a_{t,s})$  discretization errors. After all we obtain the following expression for a general form of the quark action on the anisotropic lat-

tice:

$$S_q = a_t a_s^3 \sum_x \bar{q}(x) [\gamma_0 D_0 + \nu \sum_i \gamma_i D_i + m_0 - a_t r / 2 (D_0^2 + \eta \sum_i D_i^2) - a_t i g r / 4 \{ c_B \eta \sum_{ij} \sigma_{ij} F_{ij} + c_E (1 + \eta) \sum_i \sigma_{0i} F_{0i} \}] q(x), \quad (9)$$

where the Wilson parameter  $r$  is taken arbitrary.

We can determine the four parameters  $\nu$ ,  $\eta$ ,  $c_E$  and  $c_B$  at the tree-level from the on-shell quark-quark scattering amplitude. The improvement condition is that the parameters should be chosen to remove  $O(a_t)$  cutoff effects from the scattering amplitude at the on-shell point. This is an extension of the previous work[7] done on the isotropic lattice to the anisotropic one. Comparing  $\bar{u}(p') \gamma_\mu u(p)$  ( $p' \neq p$ ) on the lattice with the continuum expression, we obtain  $\nu = \eta = c_E = c_B = 1$ . On the other hand, the  $\nu$  and  $\eta$  parameters can be also determined from the quark propagator by requiring that it should reproduce the form,

$$S_q(p) = \frac{1}{Z_q} \frac{-i \sum_\mu \gamma_\mu p_\mu + m_R}{\sum_\mu p_\mu^2 + m_R^2} + (\text{no pole terms}) + O(a_{t,s}^2) \quad (10)$$

with proper choice for  $Z_q$ ,  $Z_m$ ,  $\nu$  and  $\eta$ . We find  $Z_q^{-1/2} = 1 - r m_0 a_t / 2$ ,  $Z_m = 1 - r m_0 a_t / 2$  and  $\nu = \eta = 1$  at the tree-level. It should be noticed that the Dirac spinor takes the correct relativistic form only when eq. (10) is satisfied.

Expanding one-loop contributions to the quark self-energy  $\Sigma(p, m_0)$  in terms of  $p$  and  $m_0$ , we obtain the expressions of  $Z_q^{-1/2}$ ,  $Z_m$ ,  $\nu$  and  $\eta$  up to  $O(g^2 a_t)$ . As an example, the expression of  $Z_m$  is given by

$$Z_m = \left\{ 1 + g^2 C_F / (16\pi^2) (-3 \log(\lambda^2 a_s^2) + \Delta_m^{(0)}) \right\} \times \left\{ 1 + a_t m \left( -r/2 + g^2 C_F / (16\pi^2) \Delta_m^{(1)} \right) \right\}, \quad (11)$$

where  $\lambda$  is fictitious gluon mass to regularize the infrared divergence. Figure 1 shows  $\xi$  dependence of  $\Delta_m^{(1)}$ . Here we consider  $r = 1$  and  $r = \xi$  cases with  $\eta = 1$ , which satisfy the tree-level on-shell improvement condition. We observe approximately linear dependences on  $\xi$  for  $\Delta_m^{(1)}$ , which




Figure 1.  $\xi$  dependence of  $\Delta_m^{(1)}$  in  $Z_m$ . Errors are within symbols.

tells us that  $O(g^2 a)$  contributions to  $Z_m$  are effectively of order  $g^2 m a_s = g^2 m a_t \xi$ . This leads us to conclude that  $m a_s$  errors are allowed to revive at the one-loop level. This is a reasonable conclusion in view of the on-shell improvement. As far as we know there is no symmetry on the anisotropic lattice to prohibit the higher dimensional operators multiplied by  $(m a_s)^n$ . Unless such symmetry is uncovered, the theoretical advantage of the anisotropic lattice over the isotropic one would never be confirmed.

### 3. ISOTROPIC LATTICE

Let us explain our new relativistic approach to control  $m_Q a$  errors for the heavy quarks on the isotropic lattice. The basic idea is an application of the on-shell improvement program to the finite  $m_Q a$  case. Here we assume that the leading cutoff effects are  $f_i(m_Q a, g^2, \log a) (a \Lambda_{\text{QCD}})^i$  ( $i \geq 0$ ), where  $f_i(m_Q a, g^2, \log a) (a \Lambda_{\text{QCD}})^i > f_{i+1}(m_Q a, g^2, \log a) (a \Lambda_{\text{QCD}})^{i+1}$  and  $m_Q \gg \Lambda_{\text{QCD}}$ .  $f_i(m_Q a, g^2, \log a)$  are smooth and continuous all over the range of  $m_Q a$  and have Taylor expansions at  $m_Q a = 0$  with sufficiently large convergence radii beyond  $m_Q a = 1$ . We show that our formulation removes the cutoff effects up to  $f_1(m_Q a, g^2, \log a) a \Lambda_{\text{QCD}}$  with the use of the on-shell improvement. The remaining cutoff effects  $f_2(m_Q a, g^2, \log a) (a \Lambda_{\text{QCD}})^2$  could be removed by continuum extrapolation or may be small enough

to be neglected.

In Ref.[6], we have derived the on-shell improved action from the “off-shell” improved theory, which has the axis interchange symmetry. Here we alternatively derive it without using the axis interchange symmetry from the beginning. In this case the discussion is exactly the same as for the anisotropic case in the previous section. The generic quark action at all order of  $ma$  is given by

$$S_q = a^4 \sum_x \bar{q}(x) [\gamma_0 D_0 + \nu \sum_i \gamma_i D_i + m_0 - ar_t/2D_0^2 - ar_s/2\sum_i D_i^2 - aigc_B/4\sum_{ij} \sigma_{ij} F_{ij} - aigc_E/2\sum_i \sigma_{0i} F_{0i}] q(x), \quad (12)$$

where we are allowed to choose  $r_t = 1$  and the four parameters  $\nu$ ,  $r_s$ ,  $c_E$  and  $c_B$  are to be adjusted. We remark that the parameter  $r_s$  is not redundant, while it is in Ref.[2]. As a consequence the “ $d_1$  field rotation” required in Ref.[2] is not necessary in our formulation. This difference originates from the ways to find the redundant operators: we employ the classical field equation, while Ref.[2] uses the isospectral transformation, which is originally proposed in Ref.[5]. Although it is remarked in Ref.[5] that this transformation should be local and invariant under rotations and reflections of the lattice, Ref.[2] employs the field transformation which breaks the rotational symmetry. If the transformation is restricted to be rotationally symmetric, which is actually proportional to the classical field equation, the derived conclusion is exactly the same as ours.

The leading cutoff effects  $f_0(m_Q a, g^2, \log a)$ , which reflect discretization errors of the operators  $\bar{q}(x)q(x)$ ,  $\bar{q}(x)\gamma_0 D_0 q(x)$  and  $\bar{q}(x)D_0^2 q(x)$ , are completely absorbed in  $Z_q$  and  $Z_m$ . The next leading scaling violations stem from other terms in eq.(12). Their magnitude are expected to be  $f_1^W(m_Q a, g^2, \log a)(a\Lambda_{\text{QCD}})$  for the Wilson quark action and  $f_1^{\text{SW}}(m_Q a, g^2, \log a)(a\Lambda_{\text{QCD}})$  with  $f_1^{\text{SW}}(m_Q a = 0, g^2, \log a) = 0$  for the (massless)  $O(a)$  improved SW quark action. This implies that once we fix the pole mass from some spectral quantity, the cutoff effects in other spectral quantities are at most of order  $a\Lambda_{\text{QCD}}$ , not  $(m_Q a)^n$ , for the Wilson and the (massless)  $O(a)$  improved SW quark actions. If the

four parameters in eq.(12) are properly adjusted, the remaining scaling violations are reduced to  $f_2^{\text{ours}}(m_Q a, g^2, \log a)(a\Lambda_{\text{QCD}})^2$

The four parameters  $\nu$ ,  $r_s$ ,  $c_E$  and  $c_B$  at the tree-level are determined from the on-shell quark-quark scattering amplitude in the same way as for the anisotropic case. The improvement condition that the parameters should be chosen to reproduce the continuum scattering amplitude gives  $\nu = \sinh(m_p)/m_p$ ,  $r_s = \cosh(m_p)/m_p + \sinh(m_p)(r_t - 1/m_p)/m_p$ ,  $c_E = r_t \nu$  and  $c_B = r_s$ . Among these parameters,  $\nu$  and  $r_s$  can be determined by demanding that the tree-level quark propagator should reproduce the continuum relativistic form. As expected, both methods give exactly the same values for  $\nu$  and  $r_s$ .

## 4. CONCLUSIONS

The anisotropic lattice requires the four parameters to be tuned for the  $O(a)$  improvement even in the massless case, and allows the revival of  $m_Q a_s$  errors through one-loop diagrams. As for the isotropic lattice our proposed action, which also needs the adjustment of the four parameters, can reduce the cutoff effects to be  $O((a\Lambda_{\text{QCD}})^2)$ .

This work is supported in part by the Grants-in-Aid of the Ministry of Education (Nos. 13740169, 12014202, 12640253, 12304011, 13135204).

## REFERENCES

1. G. P. Lepage *et al.*, Phys. Rev. D**46** (1992) 4052.
2. A. X. El-Khadra, A. S. Kronfeld and P. B. Mackenzie, Phys. Rev. D**55** (1997) 3933.
3. F. Karsch, Nucl. Phys. B**205** (1982) 285; G. Burgers *et al.*, *ibid.* B**304** (1988) 587.
4. K. Symanzik, Nucl. Phys. B**226** (1983) 187, 205.
5. M. Lüscher and P. Weisz, Commun. Math. Phys. **97** (1985) 59.
6. S. Aoki, Y. Kuramashi and S. Tominaga, hep-lat/0107009.
7. R. Wohlert, DESY preprint 87-069 (1987), unpublished.